

JME 2024 - Paimboeuf
99 espèces trouvées

| G_SG_VAL | ES_VAL | SP_VAL | AUT_VAL | FAMILLE |
|---------------|-----------------|-----------------|----------------------|---------|
| AGARICUS | BERNARDII | | (Quélet) Quélet | 6B1 |
| AGARICUS | XANTHODERMUS | | Genevier | 6B1 |
| AMANITA | ALSEIDES | | Hanss | 6B3 |
| AMANITA | JUNQUILLEA | | Quélet | 6B3 |
| AMANITA | RUBESCENS | | Pers. | 6B3 |
| AMANITA | SIMULANS | | Contu | 6B3 |
| BAEOSPORA | MYOSURA | | (Fr.) Singer | 3E |
| BOLBITIUS | TITUBANS | var. VITELLINUS | (Pers.) | 5D |
| BOVISTA | PLUMBEA | | Pers. | 11B |
| CALVATIA | EXCIPULIFORMIS | | (Scopoli) Perdeck | 11B |
| CHROOGOMPHUS | RUTILUS | | (Schaeff.) Miller | 1B |
| CLATHRUS | RUBER | | Pers. | 7B |
| CLAVULINA | CINEREA | | (Bulliard) Schröter | 15 |
| CLAVULINA | RUGOSA | | (Bulliard) Schröter | 15 |
| COPRINELLUS | MICACEUS | | (Bulliard) Vilgalys, | 6A2 |
| COPRINUS | LEVISTICOLENS | | E.Ludwig et P.Roux | 6A2 |
| CRINIPELLIS | SCABELLA | | (Albertini et | 3E |
| CYSTODERMA | AMIANTHINUM | | (Scopoli) Fayod | 3D |
| GANODERMA | ADSPERSUM | | (Schulzer) Donk | 18 |
| GEASTRUM | LAGENIFORME | | Vittadini | 11A |
| GEASTRUM | MORGANII | | Lloyd | 11A |
| GYMNOPIIUS | PENETRANS | | (Fr.) Murrill | 5B |
| GYROPORUS | CASTANEUS | | (Bulliard) Quélet | 1A |
| HYDNELLUM | CONCRESCENS | | (Pers.) Banker | 14C |
| INOCYBE | GEOPHYLLA | | (Bulliard) Kummer | 5A3 |
| INOCYBE | GEOPHYLLA | var. LILACINA | (Peck) Gillet | 5A3 |
| LACCARIA | AMETHYSTINA | | M.C.Cooke | 3C1 |
| LACCARIA | LACCATA | var. PROXIMA | (Boudier) Maire | 3C1 |
| LACTARIUS | ATLANTICUS | | Bon | 2B |
| LEPIOTA | LAEVIGATA | | (Lange) Lange | 6B2 |
| LEPISTA | NUDA | | (Bulliard) | 3C1 |
| LEPISTA | SORDIDA | | (C.F.Schumacher) | 3C1 |
| LERATIOMYCES | CERES | | (M.C.Cooke et | 5C |
| LYCOPERDON | ECHINATUM | | Pers. | 11B |
| MACROCYSTIDIA | CUCUMIS | | (Pers.) Josserand | 5A2 |
| MACROTYPHULA | FILIFORMIS | | (Bulliard) | 15 |
| MARASMIUS | OREADES | | (Bolton) Fr. | 3E |
| MELANOLEUCA | EXSCISSA | | (Fr.) Singer | 3C1 |
| MELANOLEUCA | IRIS | | Kühner | 3C1 |
| MYCENA | CAPILLARIPES | | Peck | 3E |
| MYCENA | SEYNII | | Quélet | 3E |
| NEOTTIELLA | RUTILANS | | (Fr.) Dennis | 27E |
| PANAEOLUS | ATER | | (Lange) Kühner et | 5D |
| PANAEOLUS | PAPILIONACEUS | | (Bulliard) Quélet | 5D |
| PARALEPISTA | FLACCIDA | | (Sowerby) Vizzini | 3C1 |
| PAXILLUS | INVOLUTUS | | (Batsch) Fr. | 1B |
| PHALLUS | HADRIANII | | Ventenat | 7A |
| PHELLODON | NIGER | | (Fr.) Karsten | 14C |
| PLUTEUS | BRUNNEORADIATUS | | Bonnard | 4C |
| PLUTEUS | CERVINUS | | (Schaeff.) Kummer | 4C |
| RHODOCOLLYBIA | BUTYRACEA | | (Bulliard) Lennox | 3E |
| RUSSULA | AMOENOLENS | | Romagnesi | 2A |

JME 2024 - Paimboeuf
99 espèces trouvées

| G_SG_VAL | ES_VAL | SP_VAL | AUT_VAL | FAMILLE |
|--------------|---------------|--------|---------------------|---------|
| RUSSULA | CESSANS | | Pearson | 2A |
| RUSSULA | FRAGILIS | | (Pers.) Fr. | 2A |
| RUSSULA | OCHROLEUCA | | Pers. | 2A |
| RUSSULA | PARAZUREA | | Jul.Schäffer | 2A |
| RUSSULA | RECONDITA | | Melera et Ostellari | 2A |
| RUSSULA | TORULOSA | | Bresadola | 2A |
| RUSSULA | XERAMPELINA | | (Schaeff.) Fr. | 2A |
| SPARASSIS | CRISPA | | (Wulfen) Fr. | 15 |
| SUILLUS | GRANULATUS | | (L.) Roussel | 1A |
| TAPINELLA | ATROTOMENTOSA | | (Batsch) Šutara | 1B |
| THELEPHORA | TERRESTRIS | | Ehrhart | 21 |
| TRICHOLOMA | AURATUM | | Gillet | 3C2 |
| TRICHOLOMA | FOCALE | | (Fr.) Ricken | 3C2 |
| TRICHOLOMA | SULPHUREUM | | (Bulliard) Kummer | 3C2 |
| VOLVARIELLA | SPECIOSA | | (Fr.) Singer | 4C |
| XEROCOMELLUS | CHRYSETERON | | (Bulliard) Šutara | 1A |